arXiv:0705.0881v1 [gr-qc] 7 May 2007

S-BRANE SOLUTIONS WITH ORTHOGONAL INTERSECTION RULES

V.D. Ivashchuk[1]

Center for Gravitation and Fundamental Metrology, VNIIMS, 46 Ozyornaya St., Moscow 119361, Russia

Institute of Gravitation and Cosmology, Peoples' Friendship University of Russia, 6 Miklukho-Maklaya St., Moscow 117198, Russia

Abstract

A family of generalized composite intersecting S-brane solutions with orthogonal intersection rules is described.

---

[1] rusgs@phys.msu.ru

1 Introduction

This paper is devoted to a description of a family of cosmological-type solutions in the model with scalar fields and fields of forms that may be also considered as generalized S-brane solutions (see [1, 2, 3, 4], and references therein).

We deal with a model governed by the action

$$S_g = \int d^D x \sqrt{|g|} \Big\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \triangle} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)](F^a)^2 \Big\} \tag{1.1}$$

where $g = g_{MN}(x) dx^M \otimes dx^N$ is a metric, $\varphi = (\varphi^\alpha) \in \mathbb{R}^l$ is a vector of scalar fields, $(h_{\alpha\beta})$ is a constant symmetric non-degenerate $l \times l$ matrix ($l \in \mathbb{N}$), $\theta_a = \pm 1$, $F^a = dA^a = \frac{1}{n_a!} F^a_{M_1 \dots M_{n_a}} dz^{M_1} \wedge \dots \wedge dz^{M_{n_a}}$ is a n_a-form ($n_a \geq 1$), λ_a is a 1-form on $\mathbb{R}^l$: $\lambda_a(\varphi) = \lambda_{a\alpha}\varphi^\alpha$, $a \in \triangle$, $\alpha = 1, \dots, l$. In (1.1) we denote $|g| = |\det(g_{MN})|$, $(F^a)^2_g = F^a_{M_1 \dots M_{n_a}} F^a_{N_1 \dots N_{n_a}} g^{M_1 N_1} \dots g^{M_{n_a} N_{n_a}}$, $a \in \triangle$. Here $\triangle$ is some finite set. For pseudo-Euclidean metric of signature $(-, +, \dots, +)$ all $\theta_a = 1$.

2 Generalized S-brane solutions

2.1 Solutions with n Ricci-flat spaces

Let us consider a family of solutions to field equations corresponding to the action (1.1) and depending upon one variable u [3] (see also [6, 7]).

These solutions are defined on the manifold

$$M = (u_-, u_+) \times M_1 \times M_2 \times \dots \times M_n, \tag{2.1}$$

where (u_-, u_+) is an interval belonging to $\mathbb{R}$, and have the following form

$$g = \Big(\prod_{s \in S} [f_s(u)]^{2d(I_s)h_s/(D-2)} \Big) \Big\{ \exp(2c^0 u + 2\bar{c}^0) w du \otimes du + \sum_{i=1}^{n} \Big(\prod_{s \in S} [f_s(u)]^{-2h_s \delta_{iI_s}} \Big) \exp(2c^i u + 2\bar{c}^i) g^i \Big\}, \tag{2.2}$$

$$\exp(\varphi^\alpha) = \left(\prod_{s \in S} f_s^{h_s \chi_s \lambda^\alpha_{a_s}} \right) \exp(c^\alpha u + \bar{c}^\alpha), \tag{2.3}$$

$$F^a = \sum_{s \in S} \delta^a_{a_s} \mathcal{F}^s, \tag{2.4}$$

$\alpha = 1, \dots, l$; $a \in \triangle$.

In (2.2) $w = \pm 1$, $g^i = g^i_{m_i n_i}(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$ is a Ricci-flat metric on M_i, $i = 1, \dots, n$,

$$\delta_{iI} = \sum_{j \in I} \delta_{ij} \tag{2.5}$$

is the indicator of i belonging to I: $\delta_{iI} = 1$ for $i \in I$ and $\delta_{iI} = 0$ otherwise.

The p-brane set S is by definition

$$S = S_e \sqcup S_m, \quad S_v = \sqcup_{a \in \triangle} \{a\} \times \{v\} \times \Omega_{a,v}, \tag{2.6}$$

$v = e, m$ and $\Omega_{a,e}, \Omega_{a,m} \subset \Omega$, where $\Omega = \Omega(n)$ is the set of all non-empty subsets of $\{1, \dots, n\}$. Here and in what follows $\sqcup$ means the union of non-intersecting sets. Any p-brane index $s \in S$ has the form

$$s = (a_s, v_s, I_s), \tag{2.7}$$

where $a_s \in \triangle$ is colour index, $v_s = e, m$ is electro-magnetic index and the set $I_s \in \Omega_{a_s, v_s}$ describes the location of p-brane worldvolume.

The sets S_e and S_m define electric and magnetic p-branes, correspondingly. In (2.3)

$$\chi_s = +1, -1 \tag{2.8}$$

for $s \in S_e, S_m$, respectively. In (2.4) forms

$$\mathcal{F}^s = Q_s f_s^{-2} du \wedge \tau(I_s), \tag{2.9}$$

$s \in S_e$, correspond to electric p-branes and forms

$$\mathcal{F}^s = Q_s \tau(\bar{I}_s), \tag{2.10}$$

$s \in S_m$, correspond to magnetic p-branes; $Q_s \neq 0$, $s \in S$. Here and in what follows

$$\bar{I} \equiv I_0 \setminus I, \qquad I_0 = \{1, \dots, n\}. \tag{2.11}$$

All manifolds M_i are assumed to be oriented and connected and the volume d_i-forms

$$\tau_i \equiv \sqrt{|g^i(y_i)|}\, dy_i^1 \wedge \ldots \wedge dy_i^{d_i}, \tag{2.12}$$

and parameters

$$\varepsilon(i) \equiv \mathrm{sign}(\det(g^i_{m_i n_i})) = \pm 1 \tag{2.13}$$

are well-defined for all $i = 1, \ldots, n$. Here $d_i = \dim M_i$, $i = 1, \ldots, n$; $D = 1 + \sum_{i=1}^n d_i$. For any set $I = \{i_1, \ldots, i_k\} \in \Omega$, $i_1 < \ldots < i_k$, we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \ldots \wedge \tau_{i_k}, \tag{2.14}$$

$$d(I) \equiv \sum_{i \in I} d_i, \tag{2.15}$$

$$\varepsilon(I) \equiv \varepsilon(i_1) \ldots \varepsilon(i_k). \tag{2.16}$$

The parameters h_s appearing in the solution satisfy the relations

$$h_s = (B_{ss})^{-1}, \tag{2.17}$$

where

$$B_{ss'} \equiv d(I_s \cap I_{s'}) + \frac{d(I_s) d(I_{s'})}{2 - D} + \chi_s \chi_{s'} \lambda_{a_s \alpha} \lambda_{a_{s'} \beta} h^{\alpha\beta}, \tag{2.18}$$

$s, s' \in S$, with $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$. (In (2.3)) $\lambda^{\alpha}_{a_s} = h^{\alpha\beta} \lambda_{a_s \beta}$.)

Here we assume that

$$\textbf{(i)} \qquad B_{ss} \neq 0, \tag{2.19}$$

for all $s \in S$, and

$$\textbf{(ii)} \qquad B_{ss'} = 0, \tag{2.20}$$

for $s \neq s'$, i.e. canonical (orthogonal) intersection rules are satisfied.

The moduli functions read

$$f_s(u) = R_s \sinh(\sqrt{C_s}(u - u_s)), \ C_s > 0, \ h_s \varepsilon_s < 0; \tag{2.21}$$

$$R_s \sin(\sqrt{|C_s|}(u - u_s)), \ C_s < 0, \ h_s \varepsilon_s < 0; \tag{2.22}$$

$$R_s \cosh(\sqrt{C_s}(u - u_s)), \ C_s > 0, \ h_s \varepsilon_s > 0; \tag{2.23}$$

$$|Q^s| |h_s|^{-1/2} (u - u_s), \ C_s = 0, \ h_s \varepsilon_s < 0, \tag{2.24}$$

where $R_s = |Q_s| |h_s C_s|^{-1/2}$, C_s, u_s are constants, $s \in S$.

Here

$$\varepsilon_s = (-\varepsilon[g])^{(1 - \chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \tag{2.25}$$

$s \in S$, $\varepsilon[g] \equiv \mathrm{sign}(\det(g_{MN}))$. More explicitly (2.25) reads: $\varepsilon_s = \varepsilon(I_s)\theta_{a_s}$ for $v_s = e$ and $\varepsilon_s = -\varepsilon[g]\varepsilon(I_s)\theta_{a_s}$ for $v_s = m$.

Vectors $c = (c^A) = (c^i, c^\alpha)$ and $\bar{c} = (\bar{c}^A)$ obey the following constraints

$$\sum_{i \in I_s} d_i c^i - \chi_s \lambda_{a_s \alpha} c^\alpha = 0, \qquad \sum_{i \in I_s} d_i \bar{c}^i - \chi_s \lambda_{a_s \alpha} \bar{c}^\alpha = 0, \quad s \in S, \tag{2.26}$$

$$c^0 = \sum_{j=1}^{n} d_j c^j, \qquad \bar{c}^0 = \sum_{j=1}^{n} d_j \bar{c}^j, \tag{2.27}$$

$$\sum_{s \in S} C_s h_s + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^{n} d_i (c^i)^2 - \left(\sum_{i=1}^{n} d_i c^i \right)^2 = 0. \tag{2.28}$$

Here we identify notations for g^i and $\hat{g}^i$, where $\hat{g}^i = p_i^* g^i$ is the pullback of the metric g^i to the manifold M by the canonical projection: $p_i : M \to M_i$, $i = 1, \dots, n$. An analogous agreement will be also kept for volume forms etc.

Due to (2.9) and (2.10), the dimension of p-brane worldvolume $d(I_s)$ is defined by

$$d(I_s) = n_{a_s} - 1, \quad d(I_s) = D - n_{a_s} - 1, \tag{2.29}$$

for $s \in S_e, S_m$, respectively. For a p-brane we have $p = p_s = d(I_s) - 1$.

2.2 Solutions with one curved Einstein space and $n-1$ Ricci-flat spaces

The cosmological solution with Ricci-flat spaces may be also modified to the following case:

$$\mathrm{Ric}[g^1] = \xi_1 g^1, \ \xi_1 \neq 0, \qquad \mathrm{Ric}[g^i] = 0, \ i > 1, \tag{2.30}$$

i.e. the first space (M_1, g^1) is Einstein space of non-zero scalar curvature and other spaces (M_i, g^i) are Ricci-flat and

$$1 \notin I_s, \quad \forall s \in S, \tag{2.31}$$

i.e. all "brane" submanifolds do not contain M_1.

In this case the exact solution may be obtained by a little modifications of the solutions from the previous subsection. The metric reads as follows [5]

$$g = \Bigl(\prod_{s \in S} [f_s(u)]^{2d(I_s)h_s/(D-2)} \Bigr) \Bigl\{ [f_1(u)]^{2d_1/(1-d_1)} \exp(2c^1 u + 2\bar{c}^1) \times \tag{2.32}$$

$$\times [w du \otimes du + f_1^2(u) g^1] + \sum_{i=2}^{n} \Bigl(\prod_{s \in S} [f_s(u)]^{-2h_s \delta_{iI_s}} \Bigr) \exp(2c^i u + 2\bar{c}^i) g^i \Bigr\}.$$

where

$$\begin{aligned} f_1(u) = & R\sinh(\sqrt{C_1}(u-u_1)),\ C_1>0,\ \xi_1 w>0; && (2.33)\\ & R\sin(\sqrt{|C_1|}(u-u_1)),\ C_1<0,\ \xi_1 w>0; && (2.34)\\ & R\cosh(\sqrt{C_1}(u-u_1)),\ C_1>0,\ \xi_1 w<0; && (2.35)\\ & |\xi_1(d_1-1)|^{1/2}(u-u_1),\ C_1=0,\ \xi_1 w>0, && (2.36) \end{aligned}$$

u_1 and C_1 are constants, $R = |\xi_1(d_1-1)/C_1|^{1/2}$, and

$$\sum_{s \in S} C_s h_s + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^{n} d_i (c^i)^2 - \left(\sum_{i=1}^{n} d_i c^i \right)^2 = C_1 \frac{d_1}{d_1-1}. \qquad (2.37)$$

Now, vectors $c = (c^A)$ and $\bar{c} = (\bar{c}^A)$ satisfy also additional constraints

$$c^1 = \sum_{j=1}^{n} d_j c^j, \qquad \bar{c}^1 = \sum_{j=1}^{n} d_j \bar{c}^j. \qquad (2.38)$$

All other relations from the previous subsection are unchanged.

Restrictions on p-brane configurations. The solutions presented above are valid if two restrictions on the sets of composite p-branes are satisfied [6]. These restrictions guarantee the block-diagonal form of the energy-momentum tensor and the existence of the sigma-model representation (without additional constraints) [9].

The first restriction reads

$$(\mathbf{R1}) \quad d(I \cap J) \leq d(I) - 2, \qquad (2.39)$$

for any $I, J \in \Omega_{a,v}$, $a \in \triangle$, $v = e, m$ (here $d(I) = d(J)$).

The second restriction is following one

$$(\mathbf{R2}) \quad d(I \cap J) \neq 0, \qquad (2.40)$$

for $I \in \Omega_{a,e}$ and $J \in \Omega_{a,m}$, $a \in \triangle$.

3 Scalar-vacuum solutions

Here we consider as a special case the scalar-vacuum solutions when all charges Q_s are zero.

3.1 Solution with Ricci-flat spaces

The scalar vacuum analogue of the solution from subsection 2.1 reads

$$g = \exp(2c^0 u + 2\bar{c}^0) w du \otimes du + \sum_{i=1}^{n} \exp(2c^i u + 2\bar{c}^i) g^i, \qquad (3.1)$$

$$\varphi^\alpha = c^\alpha u + \bar{c}^\alpha, \qquad (3.2)$$

$\alpha = 1, \ldots, l$; where c^0 and $\bar{c}^0$ obey to (2.27) and

$$h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^{n} d_i (c^i)^2 - \left(\sum_{i=1}^{n} d_i c^i \right)^2 = 0. \qquad (3.3)$$

A special cases of this Kasner-like solution were considered in [10] (without scalar fields) and in [11] (with one scalar field).

3.2 Solutions with one Einstein space of non-zero curvature

The scalar vacuum analogue of the solution from subsection 2.2 is the following one

$$g = \exp(2c^1 u + 2\bar{c}^1)[w du \otimes du + f_1^2(u) g^1] + \sum_{i=2}^{n} \exp(2c^i u + 2\bar{c}^i) g^i, \qquad (3.4)$$

where $f_1(u)$ is defined in (2.33)-(2.36) and

$$h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^{n} d_i (c^i)^2 - \left(\sum_{i=1}^{n} d_i c^i \right)^2 = C_1 \frac{d_1}{d_1 - 1}. \qquad (3.5)$$

Here c^1 and $\bar{c}^1$ obey to linear constraints (2.38).

Special cases of this solution were considered in [10] (without scalar fields) and in [12, 14, 13] (for one scalar field).

4 Minisuperspace-covariant notations

Here we present for completeness minisuperspace-covariant relations for constraints, that explain the notion of ”orthogonal intersection rules”.

Let (see [5, 9])

$$(\bar{G}_{AB}) = \left(\begin{array}{cc} G_{ij} & 0 \\ 0 & h_{\alpha\beta} \end{array} \right), \qquad (\bar{G}^{AB}) = \left(\begin{array}{cc} G^{ij} & 0 \\ 0 & h^{\alpha\beta} \end{array} \right) \tag{4.1}$$

be, correspondingly, a (truncated) target space metric and inverse to it, where (see [8])

$$G_{ij} = d_i \delta_{ij} - d_i d_j, \qquad G^{ij} = \frac{\delta^{ij}}{d_i} + \frac{1}{2 - D}, \tag{4.2}$$

and

$$U^s_A c^A = \sum_{i \in I_s} d_i c^i - \chi_s \lambda_{a_s \alpha} c^\alpha, \quad (U^s_A) = (d_i \delta_{iI_s}, -\chi_s \lambda_{a_s \alpha}), \tag{4.3}$$

are co-vectors, $s = (a_s, v_s, I_s) \in S$ and $(c^A) = (c^i, c^\alpha)$.

In what follows we use a scalar product [9]

$$(U, U') = \bar{G}^{AB} U_A U'_B, \tag{4.4}$$

for $U = (U_A), U' = (U'_A) \in \mathbb{R}^N$, $N = n + l$.

The scalar products for vectors U^s were calculated in [9]

$$(U^s, U^{s'}) = B_{ss'}, \tag{4.5}$$

where $s = (a_s, v_s, I_s)$, $s' = (a_{s'}, v_{s'}, I_{s'})$ belong to S and $B_{ss'}$ are defined in (2.18). Due to relations (2.20) U^s-vectors are orthogonal, i.e.

$$(U^s, U^{s'}) = 0, \tag{4.6}$$

for $s \neq s'$.

The linear and quadratic constraints from (2.26), (2.38) and (2.37), respectively, read in minisuperspace covariant form as follows:

$$U^s_A c^A = 0, \qquad U^s_A \bar{c}^A = 0, \tag{4.7}$$

$s \in S$,

$$U^1_A c^A = 0, \qquad U^1_A \bar{c}^A = 0, \tag{4.8}$$

and

$$\sum_{s \in S} C_s h_s + \bar{G}_{AB} c^A c^B = C_1 \frac{d_1}{d_1 - 1}. \tag{4.9}$$

In (4.8)

$$U^1_A c^A = -c^1 + d_i c^i, \qquad (U^1_A) = (-\delta^1_i + d_i, 0), \tag{4.10}$$

is a co-vector corresponding to curvature term. It obeys

$$(U^1, U^1) = \frac{1}{d_1} - 1 < 0, \qquad (U^1, U^s) = 0 \tag{4.11}$$

for all $s \in S$. The last relation (4.11) follows from (2.31).

5 Conclusions

In this paper we overviewed cosmological-type solutions with composite intersecting p-branes from [5, 3] obeying to "orthogonal" intersection rules. Our approach gives a rather systematic way to derivation of intersection rules using the integrability conditions for Toda-like systems [5, 6, 7]. In this approach intersection rules have a minisuperspace covariant form, i.e. they are formulated in terms of scalar products of brane U-vectors and are classified by Cartan matricies of (semisimple) Lie algebras. The intersection rules considered in this paper correspond to the Lie algebra $A_1 + \ldots + A_1$. Our solutions contain as special cases certain classes of S-brane solutions (e.g. from [1, 2, 4]).

Acknowledgments

This work was supported in part by the Russian Ministry of Science and Technology, Russian Foundation for Basic Research (RFFI-01-02-17312-a) and Project DFG (436 RUS 113/678/0-1(R)).

References

[1] M. Gutperle and A. Strominger, Spacelike branes, *JHEP* **0204**, 018 (2002); hep-th/0202210.

[2] C.M. Chen, D.M. Gal'tsov and M. Gutperle, S-brane solutions in supergravity theories; *Phys. Rev.* **D 66**, 024043 (2002); hep-th/0204071.

[3] V.D. Ivashchuk, Composite S-brane solutions related to Toda-type systems, *Class. Quantum Grav.* **20**, 261-276 (2003); hep-th/0208101.

[4] N. Ohta, Intersection rules for S-branes, *Phys. Lett.* **B 558**, 213 (2003); hep-th/0301095.

[5] V.D. Ivashchuk and V.N. Melnikov, Multidimensional classical and quantum cosmology with intersecting p-branes, *J. Math. Phys.*, **39**, 2866-2889 (1998); hep-th/9708157.

[6] V.D. Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.*, **41** (1) 444-460 (2000); hep-th/9907019.

[7] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.* **18**, R82-R157 (2001); hep-th/0110274.

[8] V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, On Wheeler-DeWitt Equation in Multidimensional Cosmology, *Nuovo Cimento* **B 104**, 575 (1989).

[9] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the Generalized Composite p-branes, hep-th/9705036; *Class. Quantum Grav.* **14**, 3001-3029 (1997); Corrigenda *ibid.* **15** (12), 3941 (1998).

[10] V.D. Ivashchuk, Multidimensional Cosmology and Toda-like Systems, *Phys. Lett.* , **A 170**, 16-20 (1992).

[11] U. Bleyer and A. Zhuk, Kasner-like, inflationary and steady-state solutions in multidimensional cosmology, *Astron. Nachrichten*, **317**, 161-173 (1996).

[12] U. Bleyer and A. Zhuk, Multidimensional integrable cosmological models with negative external curvature, *Grav. Cosmol.*, **1**, 106-118 (1995).

[13] U. Bleyer and A. Zhuk, Multidimensional integrable cosmological models with positive external space curvature, *Grav. Cosmol.* **1**, 37-45 (1995).

[14] V.D. Ivashchuk and V.N.Melnikov, Multidimensional classical and quantum cosmology with perfect fluid, *Grav. Cosmol.*, **1**, 133-148 (1995); hep-th/9503223.